

3rd Targeted Protein Degradation Conference in Japan

Date

29-31st July 2025

Venue

Shonan iPark (Fujisawa, Kanagawa)

9:30 —● Registration
Networking & Coffee

10:30 —● Opening Remarks
Katsuhiro Uto, CEO, FIMECS

Morning Session, Chair: Katsuhiro Uto

10:35 —● Keynote Presentation
Title: TBD
Gwenn Hansen, CSO, Nurix Therapeutics, Inc.

11:20 —● Targeted Degradation of TRK for Cancer Therapy and Pain Management
Jialiang Wang, Executive VP of Operation, Cullgen, Inc.

- Targeted degradation of TRK overcomes resistance to kinase inhibitors
- TRK degrader is well tolerated in cancer patients
- TRK degradation exhibits significant analgesic activities in multiple preclinical pain models

**11:45 —● Discovery and Clinical Development of Targeted Protein Degraders and Degrad-
Antibody Conjugates (DACs)**
Koichi Ito, Senior Director Biology, Prelude Therapeutics

- Lessons learned from the pre-clinical discovery, translational research and development of our VHL-based IV and CRBN-based oral clinical SMARCA2 selective degraders
- Degrad-er-Antibody Conjugate (DAC) strategy to enhance drug exposure and improve the therapeutic window in pre-clinical models

12:10 —● Luncheon Sponsor Showcase and Coffee Break (Lunch box will be served)

Afternoon Session 1, Chair: Ian Churcher

13:20 —● Development of highly active PROTACs and research on expanding PROTAC molecular modalities
Hidetomo Yokoo, Chief, Division of Organic Chemistry, National Institute of Health and Science (NIHS)

- Development of peptide-based PROTAC
- Development of Nanoparticle-based PROTAC
- in silico design of PROTAC
- Comparison of several E3 ligases in the developemnt of PROTAC

13:45 —● Advancing Degrad-er Discovery and Expanding the Landscape of E3 Ligase by RaPPIDS
Kanae Gamo, CSO, FIMECS, Inc.

- Leveraging the RaPPIDS™ platform to accelerate degrad-er discovery via high-throughput synthesis and phenotypic screening
- Identification and characterization of proprietary ligands that engage novel E3 ligases
- Enabling expansion of E3 ligase usage to expand degrad-er space and accelerate pipeline progression

14:10 —● Characterization of PROTAC specificity and endogenous protein interactomes using ProtacID
Suman Shrestha, Postdoctoral Researcher, Princess Margaret Cancer Centre

- We developed ProtacID, a flexible BioID (proximity-dependent biotinylation)-based approach to identify PROTAC-proximal proteins in living cells.
- ProtacID analysis of VHL- and CRBN- recruiting PROTACs targeting a number of different proteins (localized to chromatin or cellular membranes) demonstrates how this technique can be used to validate PROTAC degradation targets and identify non-productive (i.e. nondegraded) PROTAC-interacting proteins, addressing a critical need in the field of PROTAC development.
- We also demonstrate that ProtacID can be used to characterize native, endogenous multiprotein complexes without the use of antibodies, or modification of the protein of interest with epitope tags or biotin ligase tagging.

14:35 —● Networking & Coffee Break

15:20 —● Protein Knockdown Mediated by Auxin-Inducible Degron (AID) Technology

Masato Kanemaki, Professor, Department of Chromosome Science, National Institute of Genetics

- Overview and history of auxin-inducible degron (AID) technology
- AID2-mediated protein knockdown in cell culture and animals
- Single-chain antibody AID2 (scAb-AID2) for degrading endogenous proteins without degron tag

15:45 —● Linker free PROTACs

Hai Rao, Professor, Southern University of Science and Technology

- A set of miniPROTACs with short and interchangeable degrons
- 19 amino acids can be adapted possibly without the need of linkers
- 6 ubiquitin E3 enzymes may be employed
- Excellent oral bioavailability

16:10 —● Induction of lysosomal degradation by IAP-based PROTACs, SNIPERs

Mikihiko Naito, Professor, Graduate School of Pharmaceutical Sciences, University of Tokyo

- SNIPERs recruit IAP ubiquitin ligases for protein degradation
- Identification of a target protein degraded exclusively by lysosome, not by proteasome, upon SNIPER treatment
- Mechanism of the lysosomal degradation induced by SNIPERs

16:35 —● Partner's Presentation: WuXi AppTec**17:05 —● Short Break****17:30 —● Reception Party at the Venue (Open to All Participants)****19:00 —● Venue Closing**

July 30th

Day
2**9:00 —● Registration**

Networking & Coffee

Morning Session 1, Chair: Kanae Gamo**9:30 —● Brain-penetrant molecular glue degraders targeting ALK via a novel degron: A potential therapeutic approach for ALK-positive NSCLC**

Zheng Wang, Principal Scientist, Bristol Myers Squibb

- Molecular glue degraders for ALK identified
- Novel non-loop degron distal to ATP-binding pocket
- Degraders effective on all ALK fusions and TKI-resistant ALK mutants
- Improved brain penetration of ALK degrader to treat brain metastasis

9:55 —● Oral Therapies for Immunology: Degrading IRF5, a Highly Credentialed, Historically Undrugged Transcription Factor

Veronica Campbell, Senior Director, Immunology, Kymera Therapeutics

- Building a pipeline of oral therapies for well validated targets in immunology
- Biology of IRF5 and drugging the undrugged
- KT-579, a potent, selective, oral IRF5 degrader, preclinical profile in human primary cells, patient derived cells, and in vivo disease models

10:20 —● Broad-Spectrum Efficacy of CEACAM6-Targeted ADC with BET Protein Degradar in Solid Cancer Models

Hiroyuki Kogai, Scientist, Eisai Co., Ltd.

- Developing a novel ADC targeting CEACAM6 with BET protein degrader
- Demonstrating potent and broad spectrum efficacy in pancreatic cancer, colorectal cancer, breast cancer, and lung cancer models
- Discussing combination therapy effects with immune checkpoint inhibitors

10:45 —● Coffee Break

Morning Session 2, Chair: Masato Kanemaki

11:05 —● **Bifunctional Molecules to Break Antimicrobial Resistance**

Emilia Taylor, PhD Researcher, The University of Oxford

- Antimicrobial resistance (AMR) represents a critical global health challenge, necessitating innovative therapeutic strategies with novel mechanisms of action.
- Identification of resistance enzymes in bacteria that are susceptible to targeted protein degradation via recruitment or activation of bacterial proteases, alongside the development of biochemical assays to quantify degradation potency.
- Evaluation of active and inactive BacPROTACs, demonstrating selective degradation of bacterial resistance enzymes *in vitro*

11:30 —● **Development of anti-virulence therapy against bacterial infections.**

Minsoo Kim, Associate Professor, Kyoto University

- Pathogenic bacteria utilize conserved bacterial ubiquitin ligases to degrade host proteins and evade immune responses.
- We developed a novel anti-virulence drug using PROTAC technology to selectively degrade bacterial virulence factors.
- Our PROTAC effectively inhibited bacterial growth and reduced host cell death in Salmonella-infected cells.
- This approach offers a new therapeutic modality that disables bacterial virulence with long-lasting effects, fewer side effects, and reducing resistance pressure.

11:55 —● **Partner's Presentation: FUJIFILM**

12:05 —● Luncheon Sponsor Showcase and Coffee Break (Lunch box will be served)

Afternoon Session 1, Chair: Jo Hyunsun

13:15 —● **Title: TBD**

Jose Santos, Vice President, Nurix Therapeutics

13:40 —● **MicDrop - Cellular DEL Screen in Droplets for TPD and beyond**

Ken Yamada, Associate Director, Novartis Biomedical Research

- Background on CRBN glue discovery
- Overview of cellular DEL in droplets
- Validation screen with CRBN glues for cellular IKZF3 degradation
- Screen with VHL glues for CDO1 degraders and CDO1-VHL glues

14:05 —● **New approaches to PROTACs and Molecular Glues discovery combining AI, automated synthesis and innovative biology platform**

Yann Gaston-Mathé, CEO and Co-Founder, Iktos

- Acceleration of the DMTA (Design–Make–Test–Analyze) cycle through the integration of AI-driven molecular design and robotics-enabled synthesis and screening.
 - MT-Bench: a patented, automated in-cellular imaging platform for high-throughput screening of modulators targeting protein–protein interactions (PPIs).
 - Case studies illustrating the discovery of PROTACs and molecular glues enabled by MT-Bench technology.
 - Iktos's strategy for discovering novel targeted protein degraders by combining MT-Bench with 3D GenAI technologies for AI-driven drug design.
-

14:30 —● Coffee Break

14:50 —● High-throughput proteomic screening to identify and validate novel degrader molecules and targets

Henrik Daub, Founder and CSO, NEOsphere Biotechnologies

- High-throughput proteomics rapidly screens entire libraries to identify innovative degrader compounds and disease-relevant targets at scale
- Unbiased SAR analysis and proteome-wide assessment of degrader selectivity, potency, kinetics, and efficacy support lead optimization
- Global ubiquitinomics and high-throughput proximity labeling provide insights into compound-induced ubiquitination and ternary complex formation
- The neoVERSE data analysis suite offers comprehensive analysis of large-scale proteomics data, transforming them into actionable insights for drug discovery

15:15 —● Generative AI in Target Protein Degradation: What It Can Do Now and Its Future Perspective

Tasuku Ishida, Lead Scientist, Elix, Inc.

- Overview of how AI is contributing to accelerate drug discovery
- The potential of generative AI in developing molecular glue degraders
- PROTACs and generative AI: how AI can drive PROTAC research

15:40 —● Sponsor Presentation: CellFree Sciences

15:50 —● Networking & Coffee Break

Evening Session, Chair: Gwenn Hansen**16:30 —● Strategies for non-clinical toxicity assessment of targeted protein degraders and their challenges**

Masako Imaoka, Senior Director, Daiichi Sankyo Co., Ltd.

- TPDs demonstrate powerful and sustained pharmacological effects; however, there are concerns regarding the potential for unintended protein degradation, which could lead to prolonged toxicities.
- Off-target protein degradation poses a significant risk because it may go undetected in animal studies. Establishing evaluation methods is a major challenge.
- The presentation discusses the current state of preclinical safety evaluation for TPDs and the challenges faced in strategic planning, comparing them to traditional small molecules.

16:55 —● Ten years of therapeutic TPD. What have we learnt and where next?

Ian Churcher, Chief Executive Officer, Janus Drug Discovery Consulting Ltd

- Provide a perspective on global progress in the TPD field since 2015 from discovery to clinical studies
- Reflect on where TPD has made the biggest impact, what has been achieved and what we have learnt
- Assess how TPD concepts have allowed the expansion of the wider Induced-Proximity field
- Looking ahead as to what we can expect the next generations of TPD & induced-proximity approaches to deliver

**17:20 —● Panel Discussion, Facilitator: Katsuhiko Uto
Rohan Beckwith, Gwenn Hansen, Ian Churcher and Ken Yamada
Contents of the panel discussion will be described here.****18:05 —● End of Day 2**

9:00 —● Registration
Networking & Coffee

Morning Session 1, Chair: Jose Santos

9:30 —● MrTAC opens enables induced lysosomal proteolysis of the cytosolic proteome

Lauren Veronica Albercht, Assistant Professor, University of California Irvine

- Arginine methylation is an endogenous degradative modification for lysosomal proteolysis during growth factor signaling. Here, we develop MrTAC (methylarginine targeting chimera), a small molecule that induces proximity of target proteins with methyltransferases for targeted lysosomal proteolysis.
- We confirm proof-of-concept using a dual covalent molecule of protein arginine methyltransferase 1 (PRMT1) and glycogen synthase kinase 3 (GSK3). Lysosomal proteolysis was induced rapidly, within 60 minutes, and could be sustained over multiple days.
- Mechanistically, we validate specificity to the lysosome and confirm that methylation is required for degradation.
- Our latest work validates MrTAC with endogenous neo-substrate degradation. Together, MrTAC is a potent degrader modality that exploits an entirely novel proteolytic pathway.

9:55 —● Combining The Power of Protein Degraders With the Precision of Antibodies for Synergistic Potential Using Degradant-Antibody Conjugates (DACs)

Sang Hyun Lee, Vice President, Orum Therapeutics

- Delving into the mechanism and design of an antibody degrader conjugate
- Evaluating anti-tumor activity in preclinical models
- Discussing the rationale for the chosen format and value proposition over traditional approaches

10:20 —● Title: TBD

Chinatsu Sakata-Sakurai, Head of Oncology Research, Astellas Pharma

10:45 —● Networking & Coffee Break

Morning Session 2, Chair: Jialiang Wang

11:15 —● Innovative drug discovery using protein degrader

Hyunsun Jo, Founder and CEO, Pin Therapeutics

- Clinical development strategy with CK1 α highly-selective MGD in oncology
- DAC payload strategy to minimize hematologic toxicity

11:40 —● A Molecular Glue Degradant of HuR to Treat Cancer and Cancer-induced Cachexia

Yong Cang, CSO and Co-founder, Degron Therapeutics

- Human antigen R (HuR), encoded by Elavl1, is an RNA binding protein driving tumor development and metastasis but undruggable by traditional approaches;
- Molecular glue degraders of HuR was identified from customized chemical library of DegronTx and exhibited superb activity against BRAF-mutant tumors.
- Genetic and pharmacological depletion of HuR rescued cancer-induced cachexia in mice.
- FDA has cleared DEG6498, an oral HuR degrader, for clinical development to treat solid tumors including hepatocellular carcinoma and BRAF-mutant colorectal tumor.

12:05 —● Expanding the Druggable Genome with Molecular Glues

Rohan Beckwith, SVP Chemistry, Neomorph

- Molecular glues can be utilized to target proteins lacking ligandable sites
- Neomorph have developed proprietary approaches towards rational, prospective design of molecular glues beyond CRBN
- Neomorph has discovered unique binders to new E3 ligases and leveraged these systems against novel neosubstrates with glue optimization through rational design
- Proprietary novel degrons drives expansion of the druggable genome

12:30 —● Closing Remark

Kanae Gamo, CSO, FIMECS



Always wear your name badge to access the conference space



Wi-Fi

SSID : iParkGuestNet

PW : iPark_201804




No Photo

No Record

No Post of Presentation

No entry except for the conference space (orange)



These rooms are available for business meetings on a first-come, first-served basis

Transportation

To
Shonan iPark

	From Fujisawa North Exit	From Ofuna
7	0 9 19 26 33 43 53	0 13 25 34 44 54
8	3 13 23 30 37 45 56	4 11 18 25 32 39 46 54
9	4 24 46	12 35

	To Fujisawa	To Ofuna
17	3 14 18* 24 35* 40 54 57*	5 11 26 41 56
18	9 18* 27 35* 42 57	5 11 26 41 56
19	13 26 41 55	9 24 39 54
20	14 41	9 39 54

*for express



From
Shonan iPark



0466-22-2191 (Enoshima Taxi)
0467-46-5115 (Ofuna Chuo Koutsu)

Platinum Partner



药明康德
WuXi AppTec

Silver Partner



FUJIFILM

Bronze Partner



MediRama
CONNECTING SCIENCE to LIFE



Promega



MOLSIS
Molecular Simulation and Informatics Systems



Schrödinger
シュレーディンガー株式会社

NARD institute, ltd.



KIKO-TECH



Patcore



FIMECS



iPark
Shonan